Pregnancy and breast feeding after computed tomography (CT) imaging



Pregnancy and computed tomography

X-ray contrast dye is used to show up your blood vessels on a CT scan. By using the dye, the doctor has more detailed pictures.

The x-ray contrast injection is safe to be given to pregnant women and is not radioactive*.

After you have been given the x-ray contrast, the **thyroid** function of your baby should be checked in the first week after birth*.

If you have any further questions please don't hesitate to speak to your doctor.

Breast feeding and computed tomography

After you have had the CT scan with the x-ray contrast, you can continue breast feeding as usual.



*ESUR Guidelines on Contrast Media. European Society of Urogenital Radiology. Version 7.0 2008.



Produced with support from Sheffield Hospitals Charity – working together we can help local patients feel even better

To donate visit www.sheffieldhospitalscharity.org.uk Registered Charity No 1169762



Alternative formats can be available on request. Email: alternativeformats@sth.nhs.uk

© Sheffield Teaching Hospitals NHS Foundation Trust 2017

Re-use of all or any part of this document is governed by copyright and the "Re-use of Public Sector Information Regulations 2005" SI 2005 No.1515. Information on re-use can be obtained from the Information Governance Department, Sheffield Teaching Hospitals. Email infogov@sth.nhs.uk